

Castration information sheet

Following your horse's castration please monitor the wound for any excessive bleeding, swelling or tissue protruding from the wound. The odd drips of blood over the following 12 hours can be normal, but please contact the clinic if there is any more, particularly if it is dripping fast so that you cannot count the drops.

A swelling the size of an orange is acceptable, the size of a melon is unacceptable and must be seen by a vet. If there is any tissue seen to be protruding from the wound please contact a vet immediately.

It is not necessary to clean the castration site. The site should be checked for swelling twice daily for two weeks after surgery.

Turn out is recommended. For older horses walking exercise can be resumed immediately following the surgery.

Directors

D Ballard BVM&S CertEP MRCVS
I Camm BVSc CertEP MRCVS
O Crowe BVSc CertES(Orth) DipECVS MRCVS
European Specialist in Equine Surgery
R Hepburn BVSc MS(Hons) CertEM(IntMed)
DipACVIM MRCVS ACVIM & RCVS Specialist in
Equine Medicine

S Milnes BVSc CertEP MRCVS
L Paul BSc(Hons) BVM&S CertEP MRCVS
P Ravenhill BVSc CertEM(Stud Med) BAEDT MRCVS
C Shepherd BVSc MRCVS
C Wright BVSc CertVR CertEM(IntMed) MRCVS